

**Test Report**

Number: GZHH00358769

Applicant: Orcagel Ltd, Blackhouse Circle, Blackhouse Industrial Estate, Peterhead, AB42 1BN

Date: Apr 23, 2020

Sample Description:

One (1) style of submitted sample said to be :

Item Name : **Orcagel Hand Sanitiser Gel – 70% alcohol.**

Item Quantity : 5pcs.

Date Sample Received : Apr 08, 2020

Tests conducted:

As requested by the applicant, refer to attached page(s) for details.

\*\*\*\*\*

Intertek GM Testing Service Zhuhai Co. Ltd.

*Sarah Xu*

Sarah Xu  
Asst. Manager  
Healthcare and Beauty Products



Page 1 of 6



Tests Conducted

1 Disinfection Efficacy Test

**With reference to standard test method BS EN 1500:2013** Chemical disinfectants and antiseptics- Hygienic handrub -Test method and requirements (phase 2/step2)

The testing procedure is as below:

a). Preparation of the contamination fluid

Grow the E.coli in two tubes each containing 5mL of TSB for 18h to 24h at (36±1) °C. Inoculate these cultures into two bottles with 1L TSB each and incubate for 18h to 24h at (36±1) °C. The contamination fluid is test bacterial suspension.

b). Application of the contamination fluid

Preare the hands by washing for 1 min with 5mL diluted soft soap to remove natural transients. Dry them thoroughly on paper towels. Pour the contamination fluid into the container and immerse the hands up to the mid-metacarpals for 5s with fingers spread apart. Carefully allow surplus liquid to drain back into the container.

c). Allow the hands to dry in the air for 3 min, holding them in a horizontal position with the fingers spread out and rotating them to and fro to avoid the formation of droplets.

d). Prevalues

Immediately after drying, rub the fingertips(including that of the thumb) for 1min on the base of a Petri dish containing 10 mL of TSB without neutralizer, in order to assess the release of test organisms before treatment of the hands(prevalues). Use a separate Petri dish for each hand. Prepare dilutions of these sampling fluids of 10<sup>-3</sup> and 10<sup>-4</sup> in TSB. For each dilution, spread 0.1mL over the surface of a TSSA plate using glass spreaders.

e). Hygienic handrub procedure

e)(1). Immediately after sampling for the prevalues and without recontaminating the hands, the groups shall perform the handrub in accordance with either e)(2) or e)(3), as applicable.

e)(2).Reference handrub procedure R

Pour 3mL of propan-2-ol 60%(V/V) into the cupped dry hands and rub vigorously for 30s onto the skin up to the wrists in accordance with the standard handrub procedure, to ensure total coverage of the hands. Repeat with a further 3mL propan-2-ol, to give a total rubbing time of 60s.

e)(3).Handrub procedure with test product P

Pour 3mL of sample into the cupped dry hands and rub vigorously for 30s onto the skin up to the wrists in accordance with the standard handrub procedure, to ensure total coverage of the hands. Repeat with a further 3mL sample, to give a total rubbing time of 60s.

\*\*\*\*\*



## Test Report

Number: GZHH00358769

### Tests Conducted

#### f). Postvalues

Immediately afterwards, rub the fingertips and thumbtips on the base of a Petri dish containing 10mL TSB containing neutralizer for 1min. Use a separate Petri dish for each hand.

#### g). Incubation

Incubation all plates aerobically at (36±1) °C for 18h to 24h. Count the colony forming units and reincubate for a further 24h to detect any slow growing colonies.

### REQUIREMENT

Appearance product dilution: Colorless solution

Active ingredient in product: Alcohol

Application: rub in 3mL/30s, repeat once

Neutralizer: D/E Broth

Diluent (Tryptone Sodium Chloride Solution): Tryptone pancreatic digest of casein 1.0g/L, Sodium chloride 8.5g/L

Medium:

Tryptone Soya Agar (TSA)

Tryptone Soya Selective Agar (TSSA)

### TEST ORGANISMS

Escherichia coli K12 NCTC 10538 (4.9×10<sup>8</sup>cfu/mL)

\*\*\*\*\*



Tests Conducted

## TESTING RESULTS

Table 1 List of computed lg reductions and individual differences

Volunteers	Lg reduction		Difference RP-PP
	Reference handrub(RP)	Product handrub(PP)	
1	3.95	3.89	0.06
2	3.83	4.88	-1.05
3	3.80	3.47	0.33
4	3.53	3.51	0.02
5	4.41	4.38	0.03
6	4.13	2.77	1.36
7	3.57	3.68	-0.11
8	4.20	3.05	1.15
9	4.57	3.99	0.58
10	3.36	4.60	-1.24
11	2.72	3.98	-1.26
12	3.43	3.74	-0.31
13	2.74	4.95	-2.21
14	4.15	4.32	-0.17
15	3.50	2.65	0.85
16	3.54	4.43	-0.89
17	3.04	3.13	-0.09
18	3.95	3.74	0.21
19	4.10	3.10	1.00

RP: Propan-2-ol 60% V/V

PP: Test sample

A complete set of result from 19 volunteers are available and the acceptance criterias are fulfilled.

\*\*\*\*\*



**Test Report**

Number: GZHH00358769

Tests Conducted

Table 2 Sorting of individual difference and computation for Hodges-Lehmann 97.5% upper confidence limits

Computation sheet of mean pairwise differences (d1+d2)/2								
Sample No.	Sorted differences	1.36	1.15	1.00	0.85	0.58	0.33	0.21
1	1.36	1.36 <sup>1</sup>						
2	1.15	1.26 <sup>2</sup>	1.15 <sup>4</sup>					
3	1.00	1.18 <sup>3</sup>	1.08 <sup>6</sup>	1.00 <sup>7</sup>				
4	0.85	1.11 <sup>5</sup>	1.00 <sup>8</sup>	0.93 <sup>10</sup>	0.85 <sup>13</sup>			
5	0.58	0.97 <sup>9</sup>	0.87 <sup>11</sup>	0.79 <sup>15</sup>	0.72 <sup>17</sup>	0.58 <sup>31</sup>		
6	0.33	0.85 <sup>12</sup>	0.74 <sup>16</sup>	0.67 <sup>22</sup>	0.59 <sup>30</sup>	0.46 <sup>42</sup>	0.33 <sup>53</sup>	
7	0.21	0.79 <sup>14</sup>	0.68 <sup>21</sup>	0.61 <sup>26</sup>	0.53 <sup>35</sup>	0.40 <sup>48</sup>	0.27 <sup>58</sup>	
8	0.06	0.71 <sup>18</sup>	0.61 <sup>25</sup>	0.53 <sup>34</sup>	0.46 <sup>41</sup>	0.32 <sup>54</sup>		
9	0.03	0.70 <sup>19</sup>	0.59 <sup>28</sup>	0.52 <sup>37</sup>	0.44 <sup>44</sup>	0.31 <sup>55</sup>		
10	0.02	0.69 <sup>20</sup>	0.59 <sup>29</sup>	0.51 <sup>38</sup>	0.44 <sup>45</sup>	0.30 <sup>56</sup>		
11	-0.09	0.64 <sup>23</sup>	0.53 <sup>33</sup>	0.46 <sup>40</sup>	0.38 <sup>49</sup>	0.25 <sup>59</sup>		
12	-0.11	0.63 <sup>24</sup>	0.52 <sup>36</sup>	0.45 <sup>43</sup>	0.37 <sup>50</sup>	0.24 <sup>60</sup>		
13	-0.17	0.60 <sup>27</sup>	0.49 <sup>39</sup>	<u>0.42</u> <sup>47</sup>	0.34 <sup>52</sup>			
14	-0.31	0.53 <sup>32</sup>	0.42 <sup>46</sup>	0.35 <sup>51</sup>	0.27 <sup>57</sup>			
15	-0.89	0.24 <sup>60</sup>						
16	-1.05							
17	-1.24							
18	-1.26							
19	-2.21							

The differences of the individual lg Rs of RP-PP from Table 2 are sorted in the second column and in the headline according to their size in descending order.

Table 3 WILCOXON'S matched-pairs signed-ranks test

One-sided level of significance (directional test)			
Np	0.05	0.025	0.01
18	47	40	32
19	53	46	37
20	60	52	43
21	68	59	49
22	75	66	56

\*\*\*\*\*



## Test Report

Number: GZHH00358769

### Tests Conducted

From Table 3 of critical values of Wilcoxon's matched-pairs signed-ranks test the entry for n=19 and a one-sided 0.025 level of significance, the critical value of 46 is found. Hence  $c=46+1=47$ . The pairwise differences are sorted in descending order. The 47<sup>th</sup> value is 0.42. Hence the Hodges-Lehmann upper one-sided 97.5% confidence limit for the difference in lg Rs between RP and PP is 0.42, which is less than the agreed inferiority margin of 0.6. Therefore, the hypothesis of inferiority of PP is rejected and it can be concluded that the test preparation PP is non-inferior to RP.

### Conclusion:

Pass, n=19 at a one-sided 0.025 level of significance,  $c=46+1=47$ , the 47<sup>th</sup> value is 0.42, <0.6. It can be concluded that the test sample is non-inferior to Propan-2-ol 60%(V/V) under hygienic handrub procedure.

Date Sample Received: Apr 08, 2020

Testing Period: Apr 08, 2020 to May 06, 2020

\*\*\*\*\*

End of report

*This report is made solely on the basis of your instructions and/or information and materials supplied by you. It is not intended to be a recommendation for any particular course of action. Intertek does not accept a duty of care or any other responsibility to any person other than the Client in respect of this report and only accepts liability to the Client insofar as is expressly contained in the terms and conditions governing Intertek's provision of services to you. Intertek makes no warranties or representations either express or implied with respect to this report save as provided for in those terms and conditions. We have aimed to conduct the Review on a diligent and careful basis and we do not accept any liability to you for any loss arising out of or in connection with this report, in contract, tort, by statute or otherwise, except in the event of our gross negligence or wilful misconduct. This report shall not be reproduced unless with prior written approval from Intertek GM Testing Services Zhuhai Co.,Ltd.*

